



Guidelines for the Prevention and Treatment of Opportunistic Infections Among HIV-Exposed and HIV-Infected Children

Downloaded from <http://aidsinfo.nih.gov/guidelines> on 12/21/2016

Visit the AIDSinfo website to access the most up-to-date guideline.

Register for e-mail notification of guideline updates at <http://aidsinfo.nih.gov/e-news>.

***Mycobacterium avium* Complex Disease** (Last updated November 6, 2013; last reviewed November 6, 2013)

Panel's Recommendations

- Routine screening of respiratory or gastrointestinal specimens for *Mycobacterium avium* complex (MAC) microorganisms is not recommended (**BIII**), but a blood culture for MAC should be obtained to rule out disseminated disease before initiating prophylaxis (**AIII**).
- Prophylaxis with either clarithromycin or azithromycin should be offered to HIV-infected children who have advanced immunosuppression (**AII**).
 - Children aged <1 year: <750 cells/mm³
 - Children aged 1 to <2 years: <500 cells/mm³
 - Children aged 2 to <6 years: <75 cells/mm³
 - Children aged ≥6 years: <50 cells/mm³

Discontinuing Primary Prophylaxis:

- Primary prophylaxis can be discontinued in HIV-infected children aged ≥2 years receiving stable combination antiretroviral therapy (cART) for ≥6 months and experiencing sustained (>3 months) CD4 T lymphocyte (CD4) cell count recovery well above the age-specific target for initiation of prophylaxis (i.e., as in adults, >100 cells/mm³ for children aged ≥6 years (**AI**); and >200 cells/mm³ for children aged 2 to <6 years) (**BII***).

Treating Disease:

- Testing of MAC isolates for susceptibility to clarithromycin or azithromycin is recommended (**BIII**). Combination therapy with a minimum of two drugs (e.g., clarithromycin or azithromycin plus ethambutol) is recommended to prevent or delay the emergence of resistance (**AI***). Some experts use clarithromycin as the preferred first agent (**AI***), reserving azithromycin for patients with substantial intolerance to clarithromycin or when drug interactions with clarithromycin are a concern (**AII***).
- Use of rifabutin as a third drug added to the macrolide/ethambutol regimen is controversial. Some experts would add rifabutin as a third drug to the clarithromycin/ethambutol regimen, particularly in the absence of cART and in the presence of high mycobacterial counts (**CIII**); however, drug interactions should be checked carefully, and more intensive toxicity monitoring may be warranted with such combination therapy (**AIII**). Other experts recommend against using this third agent in children because of rifabutin's increased cytochrome P450 activity, which leads to increased clearance of other drugs such as protease inhibitors and non-nucleoside reverse transcriptase inhibitors, and the potential for increased toxicity associated with concomitant administration of drugs (**CIII**).
- Treatment failure is defined as the absence of clinical response and the persistence of mycobacteremia after 8 to 12 weeks of treatment. Repeat susceptibility testing of MAC isolates is recommended in this situation, and a new multidrug regimen of two or more drugs not previously used and to which the isolate is susceptible should be administered (**AIII**). Drugs that should be considered for this scenario include rifabutin, amikacin, and a quinolone.

Secondary Prophylaxis:

- Children with a history of disseminated MAC and continued immunosuppression should receive lifelong prophylaxis to prevent recurrence (**AII***). Secondary prophylaxis typically consists of continued multidrug therapy used in treatment of disease.
- Some experts recommend discontinuation of therapy in HIV-infected children who meet **all** of the following criteria:
 - Aged ≥2 years and have completed ≥12 months of treatment for MAC;
 - Remain asymptomatic for MAC;
 - Receiving stable cART (i.e., cART not requiring change for virologic or immunologic failure);
 - Have sustained (≥6 months) CD4 count recovery well above the age-specific target for initiation of primary prophylaxis (i.e., as in adults, >100 cells/mm³ for children aged ≥6 years (**AII***) and >200 cells/mm³ for children aged 2 to <6 years) (**CIII**).

Rating of Recommendations: A = Strong; B = Moderate; C = Optional

Rating of Evidence: I = One or more randomized trials *in children*[†] with clinical outcomes and/or validated endpoints; I* = One or more randomized trials *in adults* with clinical outcomes and/or validated laboratory endpoints with accompanying data *in children*[†] from one or more well-designed, nonrandomized trials or observational cohort studies with long-term clinical outcomes; II = One or more well-designed, nonrandomized trials or observational cohort studies *in children*[†] with long-term outcomes; II* = One or more well-designed, nonrandomized trials or observational studies *in adults* with long-term clinical outcomes with accompanying data *in children*[†] from one or more similar nonrandomized trials or cohort studies with clinical outcome data; III = Expert opinion

[†] Studies that include children or children/adolescents, but not studies limited to post-pubertal adolescents

Epidemiology

Mycobacterium avium complex (MAC) refers to multiple related species of nontuberculous mycobacteria (NTM) (e.g., *Mycobacterium avium*, *Mycobacterium intracellulare*, and *Mycobacterium paratuberculosis*) that are widely distributed in the environment. Recent surveillance data have shown an increasing rate of MAC infection in some regions within the United States.¹ Comprehensive guidelines on the diagnosis, prevention, and treatment of nontuberculous mycobacterial diseases were published in 2007.² These guidelines highlight the tremendous advances in laboratory methods in mycobacteriology that have expanded the number of known NTM species from 50 in 1997 to 125 in 2006. In the United States, NTM infections outnumber *Mycobacterium tuberculosis* infections and have become an important cause of pulmonary morbidity in adults.³

MAC was the second most common opportunistic infection (OI) in HIV-infected children in the United States after *Pneumocystis jirovecii* pneumonia during the era before combination antiretroviral therapy (cART), but its incidence has greatly decreased from 1.3 to 1.8 episodes per 100 person-years during that time to 0.14 to 0.2 episodes per 100 person-years during the cART era.^{4,5} MAC is ubiquitous in the environment and presumably is acquired by routine exposures through inhalation, ingestion, or inoculation.⁶ A recent population-based study in Florida of adults and children associated soil exposure, along with black race and birth outside the United States, with MAC infection.⁷ Respiratory and gastrointestinal (GI) colonization can act as portals from which infection can disseminate.⁸

MAC can appear as isolated lymphadenitis in both HIV-infected and HIV-uninfected children. Disseminated infection with MAC in pediatric HIV infection rarely occurs during the first year of life; its frequency increases with age and declining CD4 T lymphocyte (CD4) cell count, but can occur at higher CD4 counts in younger HIV-infected children than in older children or adults. It is a recognized complication of advanced immunologic deterioration among HIV-infected children.^{6,9,10}

Clinical Manifestations

Respiratory symptoms are uncommon in HIV-infected children who have disseminated MAC, and isolated pulmonary disease is rare. Early symptoms can be minimal and may precede mycobacteremia by several weeks. Symptoms commonly associated with disseminated MAC infection in children include persistent or recurrent fever, weight loss or failure to gain weight, sweats, fatigue, persistent diarrhea, and persistent or recurrent abdominal pain. Mesenteric adenitis may mimic acute appendicitis. GI symptoms can occur alone or in combination with systemic findings. Lymphadenopathy, hepatomegaly, and splenomegaly may occur. Laboratory abnormalities include anemia, leukopenia, and thrombocytopenia. Although serum chemistries are usually normal, some children may have elevated alkaline phosphatase or lactate dehydrogenase levels. These signs and symptoms also are relatively common in the absence of disseminated MAC in HIV-infected children with advanced immunosuppression.

Diagnosis

Procedures used to diagnose MAC in children are the same as those used for HIV-infected adults.¹¹ MAC is definitively diagnosed by isolation of the organism from blood or from biopsy specimens from normally sterile sites (e.g., bone marrow, lymph node). Multiple mycobacterial blood cultures over time may be required to yield a positive result. The volume of blood sent for culture also influences yield, with increased volume leading to increased yield. Use of a radiometric broth medium or lysis-centrifugation culture technique can enhance recovery of organisms from blood.

Histology demonstrating macrophage-containing acid-fast bacilli is strongly indicative of MAC infection in a patient with typical signs and symptoms, but culture is essential to differentiate nontuberculous mycobacteria from *M. tuberculosis*, to determine which nontuberculous mycobacterium is causing infection, and to perform drug-susceptibility testing. Testing of MAC isolates for susceptibility to clarithromycin or

azithromycin is recommended (**BIII**). The BACTEC™ method for radiometric susceptibility testing can be used. Resistance for clarithromycin is defined as a minimal inhibitory concentration ≥ 32 µg/mL and a minimal inhibitory concentration of ≥ 256 µg/mL for azithromycin.¹² As with tuberculosis testing, multiplex polymerase chain reaction systems have been developed for rapid identification and drug susceptibility testing, but these are currently only available in research laboratories.^{13,14}

Prevention Recommendations

Preventing Exposure

MAC is ubiquitous in the environment. Available information does not support specific recommendations regarding exposure avoidance.¹ Person-to-person transmission is not believed to be common.

Preventing First Episode of Disease

The most effective way to prevent disseminated MAC among HIV-infected children is to preserve immune function through use of effective cART. HIV-infected children who have advanced immunosuppression should be offered prophylaxis against disseminated MAC disease according to the following CD4 count thresholds (**AII**):^{15,16}

- Children aged <1 year: <750 cells/mm³
- Children aged 1 to <2 years: <500 cells/mm³
- Children aged 2 to <6 years: <75 cells/mm³
- Children aged ≥ 6 years: <50 cells/mm³

For the same reasons that clarithromycin and azithromycin are the preferred prophylactic agents for adults, either one is recommended for prophylaxis in children (**AI***); oral suspensions of both agents are commercially available in the United States. Before prophylaxis is initiated, at-risk children should be evaluated for disseminated MAC disease, including obtaining a blood culture for MAC (**AIII**). For children who cannot tolerate azithromycin or clarithromycin, rifabutin is an alternative prophylactic agent for MAC, although drug interactions and a lack of efficacy data in children limit its use (**CIII**). Combination therapy for prophylaxis has not been shown to be cost effective and increases rates of adverse events, and therefore, generally should be avoided in children (**AIII**).

Although detection of MAC in stool or the respiratory tract may precede disseminated disease, no data demonstrate a correlation between initiation of prophylaxis in patients with detectable organisms at these sites and reduced risk of developing disseminated MAC. Therefore, routine screening of respiratory or GI specimens for MAC is not recommended (**BIII**).

Discontinuing Primary Prophylaxis

On the basis of both randomized controlled trials and observational data, primary prophylaxis for MAC can be safely discontinued in HIV-infected adults who respond to cART with an increase in CD4 count.^{17,18} In a study of discontinuing OI prophylaxis among HIV-infected children whose CD4 percentages were $\geq 20\%$ for those aged >6 years and $\geq 25\%$ for those aged 2 to 6 years, 63 HIV-infected children discontinued MAC prophylaxis, and no MAC events were observed during ≥ 2 years of follow up.¹⁹ On the basis of both these findings and data from studies in adults, primary prophylaxis can be discontinued in HIV-infected children aged ≥ 2 years receiving stable cART for ≥ 6 months who experience sustained (>3 months) CD4 cell recovery well above the age-specific target for initiation of prophylaxis (i.e., as in adults, >100 cells/mm³ for children aged ≥ 6 years and >200 cells/mm³ for children aged 2 to <6 years) (**BII***). No specific recommendations exist for discontinuing MAC prophylaxis in HIV-infected children aged <2 years.

Treatment Recommendations

Treating Disease

Disseminated MAC infection should be treated in consultation with a pediatric infectious disease specialist who has expertise in pediatric HIV infection (**AIII**). Combination therapy of MAC with a minimum of 2 drugs is recommended to prevent or delay the emergence of resistance (**AI***).²⁰⁻²³ Monotherapy with a macrolide results in emergence of high-level drug resistance within weeks.²⁴

Improved immunologic status is important for controlling disseminated MAC disease; cART should be initiated in children with MAC disease who are antiretroviral (ARV) naive. However, the optimal time to start cART in this situation is unknown; many experts treat MAC with antimycobacterial therapy for 2 weeks before starting cART to try to minimize immune reconstitution inflammatory syndrome (IRIS), although whether this makes a difference is unknown (**CIII**). For children already receiving cART, it should be continued and optimized with careful attention to potential drug interactions between the ARV and antimycobacterial drugs.

Initial MAC empiric therapy should include 2 or more drugs (**AI***): clarithromycin or azithromycin plus ethambutol.²⁵ Some experts use clarithromycin as the preferred first agent (**AI***), reserving azithromycin for patients with substantial intolerance to clarithromycin or when drug interactions with clarithromycin are a concern (**AII***).²⁶ Clarithromycin levels can be increased by protease inhibitors (PI) and decreased by efavirenz, but no data are available to recommend dose adjustments for children. Azithromycin is not metabolized by the cytochrome P450 (CYP450) system; therefore, it can be used without concern for significant drug interactions with PIs and non-nucleoside reverse transcriptase inhibitors (NNRTIs).

Because a study in adults demonstrated a survival benefit with the addition of rifabutin to clarithromycin plus ethambutol, some experts would add rifabutin as a third drug to the clarithromycin/ethambutol regimen (**CIII**);²³ however, drug interactions should be checked carefully, and more intensive toxicity monitoring may be warranted if such drugs are administered concomitantly (**AIII**).²⁷ Because rifabutin increases CYP450 activity that leads to increased clearance of other drugs (e.g., PIs, NNRTIs), and toxicity might increase with concomitant administration of drugs, other experts recommend against using this third agent in children (**CIII**). Guidelines and recommendations exist for dose adjustments necessary in adults treated with rifabutin and PIs, but the absence of data in children precludes extrapolating these to HIV-infected children undergoing treatment for disseminated MAC. No pediatric formulation of rifabutin exists, but the drug can be administered mixed with foods such as applesauce. It can also be compounded in a liquid formulation by a pharmacist. Limited safety data are available from 22 HIV-infected children (median age: 9 years) who received rifabutin in combination with 2 or more other antimycobacterial drugs for treatment of MAC for 1 to 183 weeks; doses ranged from 4 mg/kg to 18.5 mg/kg, and reported adverse effects were similar to those reported in adults.²⁸ The most commonly reported dose in children has been 5 mg/kg.

Therapy is typically prolonged and depends upon response and immune reconstitution as discussed under cessation of secondary prophylaxis.

Monitoring and Adverse Events, Including IRIS

Clinically, most patients improve substantially during the first 4 to 6 weeks of therapy. A repeat blood culture for MAC should be obtained 4 to 8 weeks after initiation of antimycobacterial therapy in patients who fail to respond clinically to their initial treatment regimen. Some experts would consider a repeat blood culture for all patients with an initial positive culture, regardless of clinical response to therapy. Improvement in fever can be expected within 2 to 4 weeks after initiation of appropriate therapy. However, for those with more extensive disease or advanced immunosuppression, clinical response may be delayed, and elimination of the organism from the blood may require up to 12 weeks of effective therapy.

IRIS in patients receiving MAC therapy during cART has been reported in HIV-infected adults and children.²⁹⁻³² New onset of systemic symptoms, especially fever or abdominal pain, leukocytosis, and focal

lymphadenitis (cervical, thoracic, or abdominal) associated with preexisting but relatively asymptomatic MAC infection has occurred after the start of cART. In addition, paradoxical worsening of systemic or local symptoms of MAC may occur as the immune system is reconstituted.

In children with very low CD4 counts, the decision to begin immediate cART must take into consideration not only the urgent need for rapid immunologic improvement, but also the possibility of IRIS due to MAC. If symptoms suggestive of MAC infection are present at the time of cART initiation, the clinician should evaluate for MAC and can consider treating for MAC presumptively. cART generally should be withheld until after the first 2 weeks of antimycobacterial therapy have been completed in patients with disseminated MAC disease who have not been treated previously with or are not receiving effective cART to reduce the risk of drug interactions and complications associated with IRIS and to lower the pill burden (**CIII**). However, ART should be started as soon as possible after the first 2 weeks of antimycobacterial therapy in order to reduce the risk of developing additional AIDS-defining OIs, and to facilitate immune reconstitution and further improve the response to antimycobacterial therapy (**CIII**). Children with moderate symptoms of IRIS can be treated symptomatically with nonsteroidal anti-inflammatory drugs (NSAIDs) or, if unresponsive to NSAIDs, a short course (such as 4 weeks) of systemic corticosteroid therapy while continuing to receive cART (**CIII**).

Adverse effects from clarithromycin and azithromycin include nausea, vomiting, abdominal pain, abnormal taste, and elevations in liver transaminase levels or hypersensitivity reactions. The major toxicity associated with ethambutol is optic neuritis, with symptoms of blurry vision, central scotomata, and red-green color blindness, which usually is reversible and rare at doses of 15–25 mg/kg in children with normal renal function. Assessments of renal function, ophthalmoscopy, and (if possible) visual acuity and color vision should be performed before starting ethambutol and monitored regularly during treatment with the agent (**AIII**). Use of ethambutol in very young children whose visual acuity cannot be monitored requires careful consideration of risks and benefits.^{33,34}

Patients receiving clarithromycin plus rifabutin should be observed for the rifabutin-related development of leukopenia, uveitis, polyarthralgias, and pseudojaundice. Tiny, almost transparent, asymptomatic peripheral and central corneal deposits that do not impair vision have been observed in some HIV-infected children receiving rifabutin as part of a multidrug regimen for MAC.²⁸

Managing Treatment Failure

Treatment failure is defined as the absence of clinical response and the persistence of mycobacteremia after 8 to 12 weeks of treatment. Repeat susceptibility testing of MAC isolates is recommended in this situation, and a new multidrug regimen of 2 or more drugs not previously used and to which the isolate is susceptible should be administered (**AIII**). Drugs that should be considered for this scenario include rifabutin, amikacin, and a quinolone. Data from treating MAC in HIV-uninfected patients indicate that an injectable agent such as amikacin or streptomycin should be considered (**CIII**).^{2,3} Because dosing of these agents in children can be problematic, drug-resistant disseminated MAC should be treated with input from an expert in this disease (**AIII**). Optimization of cART is an especially important adjunct to treatment of patients in whom initial MAC therapy has failed.

Preventing Recurrence

Children with a history of disseminated MAC should be given prophylaxis to prevent recurrence (**AII***) until their immune systems are reconstituted.³⁵ Prophylaxis in this setting means continuation of multidrug therapy, because use of a single agent (clarithromycin or azithromycin) for secondary prophylaxis carries a high risk of inducing drug-resistant MAC infection.

Discontinuing Secondary Prophylaxis

On the basis of immune reconstitution data in adults^{34,36} and data in children discontinuing primary prophylaxis, some experts recommend discontinuation of secondary prophylaxis in HIV-infected children aged ≥ 2 years who have completed ≥ 12 months of treatment for MAC, remain asymptomatic for MAC, and are receiving stable cART (i.e., cART not requiring change for viral or immune failure) and who have

sustained (≥ 6 months) CD4 count recovery well above the age-specific target for initiation of primary prophylaxis (as in adults, >100 cells/mm³ for children aged ≥ 6 years (**AII***) and >200 cells/mm³ for children aged 2 to <6 years) (**CIII**). Multidrug secondary prophylaxis should be reintroduced if the CD4 count falls below the age-related threshold.

References

1. Cassidy PM, Hedberg K, Saulson A, McNelly E, Winthrop KL. Nontuberculous mycobacterial disease prevalence and risk factors: a changing epidemiology. *Clin Infect Dis*. Dec 15 2009;49(12):e124-129. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19911942>.
2. Griffith DE, Aksamit T, Brown-Elliott BA, et al. An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *Am J Respir Crit Care Med*. Feb 15 2007;175(4):367-416. Available at <http://www.ncbi.nlm.nih.gov/pubmed/17277290>.
3. Kasperbauer SH, Daley CL. Diagnosis and treatment of infections due to *Mycobacterium avium* complex. *Semin Respir Crit Care Med*. Oct 2008;29(5):569-576. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18810690>.
4. Gona P, Van Dyke RB, Williams PL, et al. Incidence of opportunistic and other infections in HIV-infected children in the HAART era. *JAMA*. Jul 19 2006;296(3):292-300. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16849662>.
5. Nesheim SR, Kapogiannis BG, Soe MM, et al. Trends in opportunistic infections in the pre- and post-highly active antiretroviral therapy eras among HIV-infected children in the Perinatal AIDS Collaborative Transmission Study, 1986-2004. *Pediatrics*. Jul 2007;120(1):100-109. Available at <http://www.ncbi.nlm.nih.gov/pubmed/17606567>.
6. Perez Mato S, Van Dyke RB. Pulmonary infections in children with HIV infection. *Semin Respir Infect*. Mar 2002;17(1):33-46. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11891517>.
7. Reed C, von Reyn CF, Chamblee S, et al. Environmental risk factors for infection with *Mycobacterium avium* complex. *Am J Epidemiol*. Jul 1 2006;164(1):32-40. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16675537>.
8. Peacock KH, Lewis L, Lavoie S. Erosive mediastinal lymphadenitis associated with *Mycobacterium avium* infection in a pediatric acquired immunodeficiency syndrome patient. *Pediatr Infect Dis J*. Jun 2000;19(6):576-578. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10877180>.
9. Hartmann P, Plum G. Immunological defense mechanisms in tuberculosis and MAC-infection. *Diagn Microbiol Infect Dis*. Jun 1999;34(2):147-152. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10354865>.
10. Keller C, Kirkpatrick S, Lee K, Paul M, Hanson IC, Gilger M. Disseminated *Mycobacterium avium* complex presenting as hematochezia in an infant with rapidly progressive acquired immunodeficiency syndrome. *Pediatr Infect Dis J*. Aug 1996;15(8):713-715. Available at <http://www.ncbi.nlm.nih.gov/pubmed/8858681>.
11. Kaplan JE, Benson C, Holmes KH, et al. Guidelines for prevention and treatment of opportunistic infections in HIV-infected adults and adolescents: recommendations from CDC, the National Institutes of Health, and the HIV Medicine Association of the Infectious Diseases Society of America. *MMWR Recomm Rep*. Apr 10 2009;58(RR-4):1-207; quiz CE201-204. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19357635>.
12. Wong DA, Yip PC, Cheung DT, Kam KM. Simple and rational approach to the identification of *Mycobacterium tuberculosis*, *Mycobacterium avium* complex species, and other commonly isolated mycobacteria. *J Clin Microbiol*. Oct 2001;39(10):3768-3771. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11574614>.
13. Shin SJ, Lee BS, Koh WJ, et al. Efficient differentiation of *Mycobacterium avium* complex species and subspecies by use of five-target multiplex PCR. *J Clin Microbiol*. Nov 2010;48(11):4057-4062. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20810779>.
14. Iamsawat S, Surawut S, Prammananan T, Leelaporn A, Jearanaisilavong J. Multiplex PCR for detection of clarithromycin resistance and simultaneous species identification of *Mycobacterium avium* complex. *Southeast Asian J Trop Med Public Health*. May 2010;41(3):590-601. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20578547>.
15. Lewis LL, Butler KM, Husson RN, et al. Defining the population of human immunodeficiency virus-infected children at risk for *Mycobacterium avium*-intracellular infection. *J Pediatr*. Nov 1992;121(5 Pt 1):677-683. Available at <http://www.ncbi.nlm.nih.gov/pubmed/1432413>.
16. Rutstein RM, Cobb P, McGowan KL, Pinto-Martin J, Starr SE. *Mycobacterium avium* intracellular complex infection in HIV-infected children. *AIDS*. Apr 1993;7(4):507-512. Available at <http://www.ncbi.nlm.nih.gov/pubmed/8099487>.

17. Currier JS, Williams PL, Koletar SL, et al, with the AIDS Clinical Trials Group 362 Study Team. Discontinuation of Mycobacterium avium complex prophylaxis in patients with antiretroviral therapy-induced increases in CD4+ cell count. A randomized, double-blind, placebo-controlled trial. *Ann Intern Med.* Oct 3 2000;133(7):493-503. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11015162>.
18. Brooks JT, Song R, Hanson DL, et al. Discontinuation of primary prophylaxis against Mycobacterium avium complex infection in HIV-infected persons receiving antiretroviral therapy: observations from a large national cohort in the United States, 1992-2002. *Clin Infect Dis.* Aug 15 2005;41(4):549-553. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16028167>.
19. Nachman S, Gona P, Dankner W, et al. The rate of serious bacterial infections among HIV-infected children with immune reconstitution who have discontinued opportunistic infection prophylaxis. *Pediatrics.* Apr 2005;115(4):e488-494. Available at <http://www.ncbi.nlm.nih.gov/pubmed/15772172>.
20. Cohn DL, Fisher EJ, Peng GT, et al. A prospective randomized trial of four three-drug regimens in the treatment of disseminated Mycobacterium avium complex disease in AIDS patients: excess mortality associated with high-dose clarithromycin. Terry Bein Community Programs for Clinical Research on AIDS. *Clin Infect Dis.* Jul 1999;29(1):125-133. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10433575>.
21. Dunne M, Fessel J, Kumar P, et al. A randomized, double-blind trial comparing azithromycin and clarithromycin in the treatment of disseminated Mycobacterium avium infection in patients with human immunodeficiency virus. *Clin Infect Dis.* Nov 2000;31(5):1245-1252. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11073759>.
22. Benson CA, Williams PL, Cohn DL, et al. Clarithromycin or rifabutin alone or in combination for primary prophylaxis of Mycobacterium avium complex disease in patients with AIDS: A randomized, double-blind, placebo-controlled trial. The AIDS Clinical Trials Group 196/Terry Bein Community Programs for Clinical Research on AIDS 009 Protocol Team. *J Infect Dis.* Apr 2000;181(4):1289-1297. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10762562>.
23. Benson CA, Williams PL, Currier JS, et al. A prospective, randomized trial examining the efficacy and safety of clarithromycin in combination with ethambutol, rifabutin, or both for the treatment of disseminated Mycobacterium avium complex disease in persons with acquired immunodeficiency syndrome. *Clin Infect Dis.* Nov 1 2003;37(9):1234-1243. Available at <http://www.ncbi.nlm.nih.gov/pubmed/14557969>.
24. Grosset J, Ji B. Prevention of the selection of clarithromycin-resistant Mycobacterium avium-intracellulare complex. *Drugs.* 1997;54 Suppl 2:23-27; discussion 28-29. Available at <http://www.ncbi.nlm.nih.gov/pubmed/9358197>.
25. Corti M, Palmero D. Mycobacterium avium complex infection in HIV/AIDS patients. *Expert Rev Anti Infect Ther.* Jun 2008;6(3):351-363. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18588499>.
26. Ward TT, Rimland D, Kauffman C, Huycke M, Evans TG, Heifets L. Randomized, open-label trial of azithromycin plus ethambutol vs. clarithromycin plus ethambutol as therapy for Mycobacterium avium complex bacteremia in patients with human immunodeficiency virus infection. Veterans Affairs HIV Research Consortium. *Clin Infect Dis.* Nov 1998;27(5):1278-1285. Available at <http://www.ncbi.nlm.nih.gov/pubmed/9827282>.
27. Powderly WG. Treatment of infection due to Mycobacterium avium complex. *Pediatr Infect Dis J.* May 1999;18(5):468-469. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10353523>.
28. Smith JA, Mueller BU, Nussenblatt RB, Whitcup SM. Corneal endothelial deposits in children positive for human immunodeficiency virus receiving rifabutin prophylaxis for Mycobacterium avium complex bacteremia. *Am J Ophthalmol.* Feb 1999;127(2):164-169. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10030558>.
29. Race EM, Adelson-Mitty J, Kriegel GR, et al. Focal mycobacterial lymphadenitis following initiation of protease-inhibitor therapy in patients with advanced HIV-1 disease. *Lancet.* Jan 24 1998;351(9098):252-255. Available at <http://www.ncbi.nlm.nih.gov/pubmed/9457095>.
30. Phillips P, Chan K, Hogg R, et al. Azithromycin prophylaxis for Mycobacterium avium complex during the era of highly active antiretroviral therapy: evaluation of a provincial program. *Clin Infect Dis.* Feb 1 2002;34(3):371-378. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11774085>.
31. Steenhoff AP, Wood SM, Shah SS, Rutstein RM. Cutaneous Mycobacterium avium complex infection as a manifestation of the immune reconstitution syndrome in a human immunodeficiency virus-infected child. *Pediatr Infect Dis J.* Aug 2007;26(8):755-757. Available at <http://www.ncbi.nlm.nih.gov/pubmed/17848894>.
32. Babiker ZO, Beeston C, Purcell J, Desai N, Ustianowski A. Mycobacterium avium complex suppurative parotitis in a patient with human immunodeficiency virus infection presenting with immune reconstitution inflammatory syndrome. *J Med Microbiol.* Nov 2010;59(Pt 11):1365-1367. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20634331>.

33. American Academy of Pediatrics. *Red Book: 2009 Report of the Committee on Infectious Diseases*. 28th ed. 28th ed. Elk Grove Village, IL 2009.
34. Lange CG, Woolley IJ, Brodt RH. Disseminated mycobacterium avium-intracellulare complex (MAC) infection in the era of effective antiretroviral therapy: is prophylaxis still indicated? *Drugs*. 2004;64(7):679-692. Available at <http://www.ncbi.nlm.nih.gov/pubmed/15025543>.
35. Aberg J, Powderly W. HIV: primary and secondary prophylaxis for opportunistic infections. *Clin Evid (Online)*. 2010;2010. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21418688>.
36. Powderly WG. Prophylaxis for opportunistic infections in an era of effective antiretroviral therapy. *Clin Infect Dis*. Aug 2000;31(2):597-601. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10987727>.

Dosing Recommendations for Prevention and Treatment of Mycobacterium avium Complex (MAC)

(page 1 of 2)

Preventive Regimen			
Indication	First Choice	Alternative	Comments/Special Issues
Primary Prophylaxis	<ul style="list-style-type: none"> • Clarithromycin 7.5 mg/kg body weight (maximum 500 mg) by mouth orally twice daily, or • Azithromycin 20 mg/kg body weight (maximum 1200 mg) orally once weekly 	<ul style="list-style-type: none"> • Azithromycin 5 mg/kg body weight (maximum 250 mg) orally once daily • Children aged >5 years: rifabutin 300 mg orally once daily with food 	<p><u>Primary Prophylaxis Indicated for Children:</u></p> <ul style="list-style-type: none"> • Aged <1 year with CD4 count <750 cells/mm³; • Aged 1 to <2 years with CD4 count <500 cells/mm³; • Aged 2 to <6 years with CD4 count <75 cells/mm³; • Aged ≥6 years with CD4 count <50 cells/mm³ <p><u>Criteria for Discontinuing Primary Prophylaxis:</u></p> <ul style="list-style-type: none"> • Do not discontinue in children age <2 years. • After ≥6 months of cART and: <ul style="list-style-type: none"> • Aged 2 to <6 years with CD4 count >200 cells/mm³ for >3 consecutive months • Aged ≥6 years with CD4 count >100 cells/mm³ for >3 consecutive months <p><u>Criteria for Restarting Primary Prophylaxis:</u></p> <ul style="list-style-type: none"> • Aged 2 to <6 years with CD4 count <200 cells/mm³ • Aged ≥6 years with CD4 count <100 cells/mm³
Secondary Prophylaxis (Chronic Suppressive Therapy)	<ul style="list-style-type: none"> • Clarithromycin 7.5 mg/kg body weight (maximum 500 mg) orally twice daily, plus • Ethambutol 15–25 mg/kg body weight (maximum 2.5 g) orally once daily, with or without food • Children aged >5 years who received rifabutin as part of initial treatment: Rifabutin 5 mg/kg body weight (maximum 300 mg) orally once daily with food 	<ul style="list-style-type: none"> • Azithromycin 5 mg/kg body weight (maximum 250 mg) orally once daily, plus • Ethambutol 15–25 mg/kg body weight (max 2.5 g) orally once daily, with or without food • Children aged >5 years who received rifabutin as part of initial treatment: Rifabutin 5 mg/kg body weight (maximum 300 mg) orally once daily with food. 	<p><u>Secondary Prophylaxis Indicated:</u></p> <ul style="list-style-type: none"> • Prior disease <p><u>Criteria for Discontinuing Secondary Prophylaxis</u></p> <p><u>Fulfillment of All of the Following Criteria:</u></p> <ul style="list-style-type: none"> • Completed ≥6 months of cART • Completed ≥12 months MAC therapy • Asymptomatic for signs and symptoms of MAC • Aged 2 to <6 years with CD4 count >200 cells/mm³ for ≥6 consecutive months • Aged ≥6 years with CD4 count >100 cells/mm³ for ≥6 consecutive months <p><u>Criteria for Restarting Secondary Prophylaxis:</u></p> <ul style="list-style-type: none"> • Aged 2 to <6 years with CD4 count <200 cells/mm³ • Aged ≥6 years with CD4 count <100 cells/mm³

Dosing Recommendations for Prevention and Treatment of Mycobacterium avium Complex (MAC)
(page 2 of 2)

Preventive Regimen			
Indication	First Choice	Alternative	Comments/Special Issues
Treatment	<p>Initial Treatment (≥ 2 Drugs):</p> <ul style="list-style-type: none"> • Clarithromycin 7.5–15 mg/kg body weight (maximum 500 mg/dose) orally twice daily plus ethambutol 15–25 mg/kg body weight (maximum 2.5 g/day) orally once daily followed by chronic suppressive therapy <p><u>For Severe Disease, Add:</u></p> <ul style="list-style-type: none"> • Rifabutin 10–20 mg/kg body weight (maximum 300 mg/day) orally once daily 	<p>If Intolerant to <u>Clarithromycin</u>:</p> <ul style="list-style-type: none"> • Azithromycin 10–12 mg/kg body weight (maximum 500 mg/day) orally once daily <p><u>If Rifabutin Cannot Be Administered and a Third Drug is Needed in Addition to a Macrolide and Ethambutol, or if a Fourth Drug is Needed in Addition to Rifabutin for Patients with More Severe Symptoms or Disseminated Disease:</u></p> <ul style="list-style-type: none"> • Ciprofloxacin 10–15 mg/kg orally twice daily (maximum 1.5 g/day), or • Levofloxacin 500 mg daily once daily, or • Amikacin 15–30 mg/kg body weight IV in 1 or 2 divided doses (maximum 1.5 g/day) 	<p>Combination therapy with a minimum of 2 drugs is recommended for at least 12 months.</p> <p>Clofazimine is associated with increased mortality in HIV-infected adults and should not be used.</p> <p>Children receiving ethambutol who are old enough to undergo routine eye testing should have monthly monitoring of visual acuity and color discrimination.</p> <p>Fluoroquinolones (e.g., ciprofloxacin and levofloxacin) are not labeled for use in children aged <18 years because of concerns regarding potential effects on cartilage; use in younger individuals requires an assessment of potential risks and benefits</p> <p>Chronic suppressive therapy (secondary prophylaxis) is recommended in children and adults following initial therapy.</p>

Key to Acronyms: cART = combination antiretroviral therapy; CD4 = CD4 T lymphocyte; MAC = *Mycobacterium avium* Complex; IV = intravenous